

Topologically coupled energy bands in molecules

Frédéric Faure*

Boris Zhilinskii†

1st February 2008

Abstract

We propose a concrete application of the Atiyah-Singer index formula in molecular physics, giving the exact number of levels in energy bands, in terms of vector bundles topology. The formation of topologically coupled bands is demonstrated. This phenomenon is expected to be present in many quantum systems.

PACS: 03.65.Vf; 02.40.-k; 31.15.Gy; 33.20.Wr

Key-Words: Born-Oppenheimer approximation; Molecular spectra; Vector bundles; Index formula; Semi-classical analysis;

1 Introduction

Molecular spectra of small molecules possess thousands of energy levels which are known with high precision and characterize the dynamical behavior of nuclei and electrons in very fine details. Thus a qualitative approach may be useful in front of the huge complexity of the spectrum [1, 2, 3]. Due to symmetries of different origins, the energy levels are often regrouped together in multiplets with quasi-degeneracies, forming so called energy bands. In terms of dynamics, the fine structure of the bands reflects a slow motion, whereas the coarser repartition of different energy bands is related to a faster motion. These two motions can be strongly coupled together.

*Laboratoire de Physique et Modélisation des Milieux Condensés (LPM2C) (Maison des Maîtres Jean Perrin, CNRS) BP 166 38042 Grenoble Cedex 9 France.
email: frederic.faure@ujf-grenoble.fr

http://lpm2c.polycnrs-gre.fr/faure/

†Université du Littoral, MREID, 145 av. M. Schumann, 59140 Dunkerque, France zhilin@univ-littoral.fr

Several typical molecular examples can be suggested to illustrate the presence of such a band structure and the evolution of this band structure with the variation of external parameters.

i) Vibrational structure of different electronic states. The existence of a fine vibrational structure associated with different electronic states is the standard consequence of the Born-Oppenheimer (adiabatic) separation of electronic and nuclear variables in molecules based on the smallness of the ratio $(m_e/M_n)^{1/4} \sim 1/10$ (here m_e and M_n are the electron mass and the typical mass of nuclei) [4].

ii) Rotational structure of different vibrational states. The rotational excitations are typically smaller than the vibrational ones. The quantum spectrum has a fine rotational band structure associated with a more crude vibrational structure. The absolute value J of the angular momentum is a strict integral of motion for an isolated molecule, and can be used as a natural parameter to analyze the evolution of rotational bands [2, 3].

iii) A more sophisticated example considered in this paper arises for systems which possess vibration modes with (quasi)-degenerate frequencies. Such a situation typically takes place in problems with symmetry. The description of the degenerate vibrations in the harmonic oscillator approximation gives an energy spectrum formed by degenerated energy levels, the so called polyads. Each polyad is characterized by the total number of vibrational quanta N , it contains. In systems where these vibrational modes are coupled with electronic states, one obtains an electronic-vibrational (vibronic) band structure. A typical question is the evolution of this vibronic band structure with N [5, 6] and its persistence (or reorganization) as a function of external physical parameters.

In all these previous examples, when an external parameter is varied, along with modifications of the internal structure of each isolated band, one observes more serious modifications of the global band structures [7, 8, 9, 10, 11]. Sometimes two bands come in contact. During this contact, a certain number of levels pass from one band to the other, resulting in a redistribution of levels between bands. These qualitative phenomena reflect band structure properties which are robust under modification of parameters, except for abrupt structure changes, and suggest a topological description. One expects a characterization of each band which would

give the precise number of levels it contains, and would predict the possible band change of structure under an external parameter variation. Such a characterization has been done for rotational bands in [12, 13]. It has been shown that an integer topological Chern index can be attributed to each individual rotational band and a generic modification of the band structures is related to the exchange of one energy level between two consecutive bands. The presence of symmetry leads to more complicated redistribution of energy levels between bands. The purpose of the present paper is to describe a new qualitative phenomenon related with the formation of topologically coupled bands.

The mathematical approach which we need to characterize these qualitative modifications is a semi-classical analysis within the Born-Oppenheimer approximation and topological analysis of vector bundles. Precise mathematical theories are developed in [14, 15, 16].

For rotational bands, the topological phenomena are quite simple due to the fact that the slow rotational motion has one degree of freedom, and takes place on a two dimensional sphere [12, 13]. We present now a more interesting example both from the physical and mathematical point of view, where the slow motion comes from molecular vibrations and has two degrees of freedom (a four dimensional phase space).

2 Semi-quantum model with CP^2 classical phase space

We first consider three vibrations with the same frequency (1:1:1 resonance) described by

$$H_{vib} = \sum_{k=1,2,3} \frac{1}{2} (p_k^2 + q_k^2) = \sum_{k=1,2,3} |Z_k|^2, \quad (1)$$

where a complex notation $Z_k = (q_k + ip_k) / \sqrt{2}$ is used. The time evolution of a point $Z = (Z_1, Z_2, Z_3)$ in phase space $\mathbb{C}^3$ is given by a phase factor: $Z(t) = \exp(-it) Z(0)$. The family of (closed) trajectories of a given energy, say $E = 1$, forms a four dimensional phase space $[4 = 6 - 1(\text{energy}) - 1(\text{time})]$ which is a reduced phase space

from the classical mechanics point of view. This space is known in mathematics as the complex projective space and noted $\mathbb{CP}^2 = P(\mathbb{C}^3)$: a point in $\mathbb{CP}^2$ corresponds to a given closed trajectory of energy E and will be noted by $[Z]$ in the sequel. The corresponding quantum Hamiltonian $\hat{H}_{vib} = \sum_{k=1,2,3} a_k^\dagger a_k + 3/2$ is obtained by replacing $Z_k, \overline{Z_k}$ by operators $a_k, a_k^\dagger$ respectively. A given energy level is thus an integer N plus $3/2$, and this level is degenerated with multiplicity

$$\mathcal{N}_0 = (N + 1)(N + 2)/2.$$

The associated eigenspace $\mathcal{H}_{polyad} = \mathbb{C}^{\mathcal{N}_0}$ is called the polyad N , and corresponds to the classical phase space $\mathbb{CP}^2$ [17].

Let us now consider a more complicated system where the previous degenerated vibrations are coupled with additional degrees of freedom. An explicit model will be given in the next section. We first discuss physical situations in molecular physics which motivate this model.

Suppose that these additional degrees of freedom have much higher energy excitations, which can be related, for example, to electronic excitation, or to another vibrational excitation with higher frequency. Due to this higher scale of energy, it is sufficient to take into account only finite (and rather low in fact) number of quantum states of this second subsystem, called the “fast” subsystem. In the absence of quasi-degeneracies in the fast subsystem one state should be sufficient, but in the case of the presence of degeneracies or quasi-degeneracies we should naturally treat together all the quasi-degenerate quantum levels.

Such a situation takes place, for example, for the dynamic Jahn-Teller effect of type $F - f$ [18], which describes the coupling of the triply degenerate vibrational modes, with a multiplet of triply degenerate electronic states. Although in the standard treatment of Jahn-Teller effect the coupling between vibrational states from different polyads is explicitly taken into account [18, 19, 20], in the limit of weak Jahn-Teller coupling, the restriction to the effective model for an isolated polyad is quite natural [21, 22]. We deal with this effective model here.

Another similar and natural example is the vibrational structure of combination bands $\nu_3 + N\nu_4$ in tetrahedral AB_4 type molecules. Due to tetrahedral symmetry both vibrational modes ν_3 and ν_4 are triply degenerate. The ν_3 frequency which is

mainly due to A-B stretching vibrations is typically higher than the ν_4 frequency, associated with bending. In the absence of resonance between ν_3 and ν_4 we can treat the combination band $\nu_3 + N\nu_4$ as formed by three bands associated with three quantum states corresponding to one quantum excitation of triply degenerate ν_3 mode. Each of these quantum states is associated with the sequence of vibrational polyads formed by ν_4 overtones.

In both cases, purely vibrational and Jahn-Teller, the mathematical model can be formalized in the same way. We consider the coupling of three quantum states with vibrational polyads formed by triply (quasi)degenerate modes. We associate the three quantum states with the high-frequency motion (it is the “fast” subsystem) and vibrational polyads with low-frequency motion (it is the “slow” subsystem). To simplify the language we will speak about “fast” subsystem as electronic and about “slow” subsystem as vibrational. The general structure of this model can be described as “semi-quantum” model. The notion “semi-quantum” used in some of earlier publications [10, 11] reflects the presence of two subsystems, one is treated as classical whereas another as purely quantum.

The simplest model assumes the degeneracy of electronic states and the degeneracy of vibrational states within polyads. Among the possible physically interesting corrections there are:

- i) vibrational couplings and anharmonicities which result in splitting of degeneracies in vibrational polyads, but make no distinction between different electronic states;
- ii) pure electronic splitting, which does not influence the internal vibrational polyad structure and can be present either (in the absence of symmetry) due to initial splitting of accidentally quasi-degenerate electronic states or produced by symmetry breaking effects like external electric or magnetic fields;
- iii) vibronic coupling between electronic and vibrational states.

We will suppose that the purely vibrational coupling is not important, because it modifies only the internal structure of energy bands. Two really different limit cases are ii) and iii) above. They correspond to situations with, on one hand, purely electronic splitting being large as compared to vibronic coupling and on the other hand with the vibronic coupling being the most essential. Naturally, this last case

corresponds to strictly degenerate (for example due to symmetry) electronic states.

The simultaneous presence of a pure electronic splitting and a vibronic coupling will be described by an explicit model in the next section, eq.(2). We will introduce an external parameter $\lambda = 0 \rightarrow 1$, which will relate the two extreme situations ii) and iii).

2.1 A matrix model

Let us describe three “electronic” states by the quantum Hilbert space $\mathcal{H}_{elec} = \mathbb{C}^3$. We suppose that vibronic (electronic-vibrational) coupling is sufficiently small as compared to vibrational frequency and can be taken into account only through interactions between polyads with the same polyad quantum number N which belong to different electronic states. This means that the total Hilbert space, we are interested in, $\mathcal{H}_{tot} = \mathcal{H}_{polyad} \otimes \mathcal{H}_{elec}$ has dimension $3\mathcal{N}_0$.

The main effect of the coupling is to remove the degeneracy of the triplets of vibrational polyads. In the “semi-quantum” description of Born-Oppenheimer approximation, when the vibrational motion is treated classically, and the electronic motion is still quantum, the coupling results in a slow precession of the orbits, that is a slow motion of the representative point in $\mathbb{C}P^2$. We will introduce the simplest form of the coupling, in a way similar to the standard spin-orbit coupling $\mathbf{S} \cdot \mathbf{L}$, but now the coupling is chosen to be invariant with respect to the $SU(3)$ symmetry group. For that purpose, we extend the $SU(3)$ symmetry of the three-dimensional harmonic oscillator (1) to the three electronic states, supposing that the space $\mathcal{H}_{elec}$ is the three-dimensional fundamental representation of $SU(3)$, and that the Hamiltonian $\hat{H}_1$ below is $SU(3)$ invariant.

In the spirit of semi-quantum approach, a classical Hamilton function is defined on phase space $\mathbb{C}P^2$ with values in the space of 3×3 Hermitian matrices:

$$(H_1([Z]))_{i,j=1,2,3} = \overline{Z_i} Z_j / \left(\sum_{k=1,2,3} |Z_k|^2 \right) = (|Z\rangle\langle Z|)_{i,j}.$$

The last expression shows that $H_1([Z])$ can be interpreted geometrically as a projector in $\mathcal{H}_{elec}$ space onto the one dimensional space, a line, defined by the value of $[Z]$. From this, the eigenvalues of $H_1([Z])$ are $E_{Line} = 1$, and $E_{Orth} = 0$ with

multiplicity two. “*Orth*” is for the two dimensional space orthogonal to the line.

Now we consider the quantum operator $\hat{H}_1$ obtained from the matrix valued function H_1 by replacing $Z_k, \overline{Z}_k$ by operators $a_k, a_k^\dagger$ respectively. We expect that $\hat{H}_1$ has two different eigenvalues close to $E_{Line} = 1$, and $E_{Orth} = 0$ (in the semi-classical limit $N \rightarrow \infty$). From its invariance under the $SU(3)$ group action, the eigenspaces of $\hat{H}_1$ are irreducible representations of this group with respective dimensions $\mathcal{N}_{Line}, \mathcal{N}_{Orth}$. The eigenspaces of $\hat{H}_1$ are obtained by Young tableau technics, see figure 1. The Weyl’s formula gives

$$\mathcal{N}_{Line} = (N + 2)(N + 3)/2,$$

$$\mathcal{N}_{Orth} = N(N + 2).$$

$$\begin{array}{ccccccc} \begin{array}{|c|c|c|c|c|c|} \hline & & & & & \\ \hline \end{array} & \otimes & \begin{array}{|c|} \hline \text{shaded} \\ \hline \end{array} & = & \begin{array}{|c|c|c|c|c|c|c|} \hline & & & & & & \text{shaded} \\ \hline \end{array} & \oplus & \begin{array}{|c|c|c|c|c|c|} \hline & & & & & \\ \hline \text{shaded} & & & & & \\ \hline \end{array} \\ \mathbf{H}_{\text{tot}} = \mathbf{H}_{\text{polyad}} & \otimes & \mathbf{H}_{\text{elec}} & = & \text{Band "Line"} & \oplus & \text{Band "Orth"} \end{array}$$

Figure 1: Spectrum of $\hat{H}_1$ in terms of $SU(3)$ representations.

The splitting of the whole system of $3(N + 1)(N + 2)/2$ energy levels into two bands is the result of the $SU(3)$ symmetry of the model Hamiltonian $\hat{H}_1$. We can now introduce the perturbation which breaks the $SU(3)$ symmetry, and analyses the persistence of the two bands and the possible evolution of the band structure. In order to see qualitative modifications of the band structure, we introduce an external parameter λ and construct a one-parameter family of Hamiltonians H_λ , $\lambda \in [0, 1]$, which relates the Hamiltonian H_1 introduced above, with a Hamiltonian H_0 which does not depend on $[Z]$ (no vibronic coupling) but possess three non-degenerate electronic states:

$$H_\lambda([Z]) = (1 - \lambda) H_0 + \lambda H_1([Z]), \quad (2)$$

$$\text{with } H_0 = \text{Diag}(-1, 0, 1).$$

For $\lambda = 0$, every eigenvalue $E = -1, 0, 1$ of the Hamiltonian $\hat{H}_0$ has multiplicity $\mathcal{N}_0$. As λ varies from 0 to 1, we get a redistribution of the energy levels from three groups of levels towards two groups, as shown on figure 2(a).

The physical interpretation of the family of Hamiltonians (2) can be done for




Figure 2: (a) The exact spectrum of $\hat{H}_\lambda$, for $N = 4$, with a transfer of $\Delta N = N + 2 = 6$ levels towards the upper group.
(b) The continuous bands spectrum in the Born-Oppenheimer approximation, interpreted as vector bundles. The Atiyah-Singer formula (3) gives the number of levels of a given band, say N_{Line} , in terms of the topology of the vector bundle V_{Line} .

both realizations mentioned above, the vibrational structure of the combination bands and the Jahn-Teller effect.

In the case of vibrational combination bands the $\hat{H}_0$ Hamiltonian corresponds to the vibrational structure formed by overtones of triply degenerate harmonic oscillator coupled with three close but non-degenerate vibrational states with higher frequencies. The $\hat{H}_1$ term introduces the interaction between low and high frequency states. It becomes leading term as soon as the splitting between high frequency states goes to zero or equivalently as soon as the coupling between low and high frequency motion becomes more important than the high frequency splitting. This means, for example, that the formal parameter λ in Eq. (2) can be associated with the polyad quantum number N which is naturally responsible for the strength of the interaction between slow and fast subsystems.

For the Jahn-Teller $F - f$ problem the Hamiltonian $\hat{H}_1$ appears in a natural way as a diagonal in polyad quantum number effective operator. At the same time the $\hat{H}_0$ limit formally corresponds to the absence of the Jahn-Teller effect and even to the absence of the three-fold degeneracy which is the origin of the Jahn-Teller interactions. To give again a physical significance to the family of Hamiltonians (2), one should introduce the reason of the splitting of electronic states. This can be formally done by adding some external magnetic field. In such a case, to approach the $\hat{H}_0$ limit, the splitting caused by magnetic field should become larger than the splitting of vibrational polyads caused by pseudo Jahn-Teller interactions, i.e. by vibronic interactions between different non-degenerate electronic states. Similar effect can be produced by any symmetry breaking effect whose influence is much more important on electronic subsystem than on vibrations.

2.2 Symmetry analysis of band decompositions

From the point of view of the Jahn-Teller dynamic effect the most interesting prediction of our topological analysis is the splitting of the whole multiplet of $3(N + 1)(N + 2)/2$ vibronic states into two groups which we will interpret later as an elementary band “Line” and a topologically coupled band “Orth.”, both with non trivial topological Chern class. From the point of view of practical applications of this statement to concrete molecules, the simple number of states is clearly

an insufficient information in order to interpret the observed energy level pattern formed by two different bands. A much more detailed and useful description of two bands can be done by indicating the symmetry of all levels forming each band. In the case of high symmetry molecules this information significantly reduces possible misinterpretation of the band structures in concrete examples. This is why, before going to more general topological characterization, we look now at the problem of the decomposition of the whole set of energy levels into different bands, from the point of view of different symmetry groups.

The highest symmetry is the $SU(3)$ symmetry which is the dynamic symmetry group of the 3-D isotropic harmonic oscillator (1). The actual invariance group of the concrete molecular problem is a finite point symmetry group G (tetrahedral $G = T_d$ or octahedral $G = O_h$ for examples mentioned in this paper). The very convenient approximate intermediate symmetry group $O(3)$ is quite useful to give additional label for groups of levels. The classification based on the chain of group $SU(3) \supset O(3) \supset G$ enables one to verify the presence of groups of levels which can be interpreted as bands.

In the case of $F_2 - f_2$ Jahn-Teller effect (the coupling of triply degenerate electronic state of F_2 symmetry with vibrational polyads formed by triply degenerate vibrations of the same symmetry for molecule with T_d symmetry group) the splitting into two bands associated with two irreducible representations of $SU(3)$ dynamical symmetry group is followed by further decomposition into irreducible representations of $O(3)$ and T_d groups. For low vibrational polyads this decomposition is given in Table 1 through the decomposition of each band into irreducible representations of the symmetry group T_d and intermediate $O(3)$ symmetry group.

The decomposition into two bands with $(N+2)(N+3)/2$ and $N(N+2)$ energy levels is imposed by the approximate dynamical $SU(3)$ symmetry which becomes an exact symmetry for the Hamiltonian (2) in the limit $\lambda = 1$. The important point now is the justification of the persistence of this decomposition found in the absence of $SU(3)$ symmetry, on the basis of topological arguments.

Table 1: Splitting of multiplets of vibrational polyads into two bands for weak $F_2 - f_2$ Jahn-Teller problem. Energy levels forming each band are listed by their symmetry labels for polyads with $N = 1, 2, 3, 4, 5$. First sub-line gives for each band the decomposition into irreducible representations of T_d symmetry group. Second sub-line gives similar decomposition in the case of $O(3)$ invariance group.

Band	Degen	$N = 1$	$N = 2$	$N = 3$
Line	$(N + 2)(N + 3)/2$	A_1, E, F_2 $0_g, 2_g$	$A_1, F_1, 2F_2$ $1_u, 3_u$	$2A_1, 2E, F_1, 2F_2$ $0_g, 2_g, 4_g$
Orth.	$N(N + 2)$	F_1 1_g	E, F_1, F_2 $1_u, 2_u$	$A_2, E, 2F_1, 2F_2$ $1_g, 2_g, 3_g$
		$N = 4$		$N = 5$
		$A_1, E, 2F_1, 4F_2$ $1_u, 3_u, 5_u$	$2A_1, A_2, 3E, 2F_1, 4F_2$ $0_g, 2_g, 4_g, 6_g$	
		$A_1, A_2, 2E, 3F_1, 3F_2$ $1_u, 2_u, 3_u, 4_u$	$A_1, A_2, 3E, 5F_1, 4F_2$ $1_g, 2_g, 3_g, 4_g, 5_g$	

3 Topological interpretation

We now proceed to the interpretation of the redistribution phenomenon observed in figure 2(a) within the Born-Oppenheimer approximation using the semi-quantum model. The matrix $H_\lambda([Z])$ has three eigenvalues $E_1(\lambda, [Z]) \leq E_2(\lambda, [Z]) \leq E_3(\lambda, [Z])$ forming three bands when $[Z] \in \mathbb{C}P^2$ varies, see figure 2(b). The vibrational motion is now treated classically, and so the fine structure of figure 2(a) (due to quantization of the vibrations), has been replaced by a continuous distribution of energies. Degeneracies between consecutive bands occur for certain values of $(\lambda, [Z])$. Between bands 2 and 3, there is a degeneracy surface in the range $1/2 < \lambda < 2/3$ ¹. For a fixed value of λ , the associated eigenspaces of the eigenvalues E_1, E_2, E_3 also depend on $[Z]$ and therefore define vector bundles² over $\mathbb{C}P^2$. The rank r of the bundle is the dimension of these eigenspaces.

The topology of a general vector bundle F over $\mathbb{C}P^2$ is characterized by its rank r and a polynomial $C(F) = 1 + Ax + Bx^2$, called the total Chern class, with integer coefficients $A, B \in \mathbb{Z}$, see [24] p.300. A rank $r = 1$ bundle has necessarily $B = 0$. The Chern class of the sum $V \oplus V'$ of two vector bundles is the product of polynomials $C(V \oplus V') = C(V) \wedge C(V')$, keeping only terms of degree not greater

¹ For a general family of Hermitian matrices, degeneracies occur with three external parameters; this is the Wigner-Von Neumann theorem, see [23], §79. Here, with the five external parameters $(\lambda, [Z])$, degeneracies are expected to hold on two-dimensional surfaces. The degeneracy surface between bands 2 and 3 is a homologically non trivial sphere S^2 in $\mathbb{R} \times \mathbb{C}P^2$.

² A vector bundle F over a space P is a family of vector space $F(p)$ which depend continuously on a parameter $p \in P$. In the Born-Oppenheimer approximation, vector bundles appear naturally: the vector space is spanned by the stationary state(s) of the fast motion, which depends on the slow classical state $p \in P$ in phase space. The topology of a vector bundle reflects the twists among the space $F(p)$ as p is varied. A vector bundle with no topology (no twist) is called trivial.

than 2.

For $\lambda < 1/2$, on figure 2(b), there are three distinct bands, and so three associated vector bundles of rank $r = 1$, noted T_1, T_2, T_3 . These bundles are trivial in the sense that for $\lambda = 0$ the eigenspaces do not depend on $[Z]$, and by continuity the topology is also trivial for any $\lambda < 1/2$. The Chern classes are $C(T_i) = 1$, with $A = B = 0$.

For $1/2 < \lambda < 2/3$, there are degeneracy surfaces, and the three previous bundles T_1, T_2, T_3 “interact” together and form a vector bundle $T = T_1 \oplus T_2 \oplus T_3$ of rank 3. This is exactly the vector space $\mathcal{H}_{elec} = \mathbb{C}^3$ which does not depend on $[Z]$, so T is also trivial, this confirms that $C(T) = 1$.

For $2/3 < \lambda < 1$, the previous bundle T decomposes into two bundles $T = V_{Line} \oplus V_{Orth}$. The upper energy band corresponds to a rank 1 vector bundle V_{Line} . By continuity, its topology can be computed at $\lambda = 1$, and corresponds to the well known non trivial canonical bundle over $\mathbb{CP}^2$ with $C(V_{Line}) = 1 + x$, see [24]. The lower band corresponds to the orthogonal bundle V_{Orth} . This is a non trivial bundle of rank $r = 2$, with $C(V_{Orth}) = 1 - x + x^2$, because $1 = C(T) = C(V_{Line}) \wedge C(V_{Orth})$. The vector bundle V_{Orth} cannot be decomposed into two rank 1 bundles just because its Chern class can not be factorized, $1 - x + x^2 \neq (1 + Ax) \wedge (1 + A'x)$ with integers A, A' . Therefore we obtain that V_{Orth} is composed of *two bands which are topologically coupled*: a continuous perturbation cannot create a gap in it. To be decoupled, they would need a topological interaction with an external third band. This interesting phenomenon seems to be new in molecular physics.

3.1 Number of energy levels in band and index theorem

The relation between the topology of a vector bundle F defined in the Born-Oppenheimer approximation, and the number of levels in the exact energy spectrum is given by the remarkable index formula of Atiyah-Singer ([24] p.330), which, in the case of the molecular model studied in this paper, reads as follows:

$$\mathcal{N}(F) = [Ch(F) \wedge Ch(Polyad) \wedge Todd(\mathbb{CP}^2)]_{/x^2}, \quad (3)$$

where $Ch(F) = r + Ax + \frac{1}{2}(A^2 - 2B)x^2$ is the Chern character for the topology of the vector bundle F . $Ch(Polyad) = \exp(Nx)$ characterizes the quantization of the polyad space³, and $Todd(\mathbb{C}P^2) = 1 + \frac{3}{2}x + x^2$ is related to the topology of the phase space $\mathbb{C}P^2$. To make calculation using the right hand side of the formula (3), one has to multiply the three polynomials, and keep the coefficient of the term x^2 . This gives

$$\mathcal{N}(F) = \frac{r}{2}N^2 + \left[\frac{3r}{2} + A\right]N + \left[r + \frac{3A}{2} + \frac{A^2}{2} - B\right].$$

It is easy to check that this formula gives the values of $\mathcal{N}_{Line}$ and $\mathcal{N}_{Orth}$ computed before. The Young tableau machinery was possible at $\lambda = 1$ thanks to a special symmetry. Here the index formula with its topological nature is much more general and is valid for any values of λ . The topology modification is related to the level exchange: in the interval $1/2 < \lambda < 2/3$ there is a transfer of $\Delta N = \mathcal{N}_{Line} - \mathcal{N}_0 = N + 2$ levels to the upper group, through the degeneracy surface⁴.

It is useful to note here that the number of quantum states for any band defined in the classical limit over $\mathbb{C}P^2$ is a quadratic function of the polyad quantum number N . The coefficients of this quadratic polynomial give information about the total Chern class and therefore the topology of the fiber bundle. In the completely classical model, when both slow and fast dynamical variables are treated through classical mechanics, a similar relation between the volume of the reduced classical phase space and Chern classes was established by Duistermaat and Heckman [26].

In comparison with the ro-vibrational model described on a S^2 phase space of dimension 2 [12], this model is much more rich due to the dimension 4 of the phase space $\mathbb{C}P^2$. The new phenomenon consists in a formation of the degeneracy surface together with the transition of a whole group of levels, instead of formation of isolated degeneracy points with transition of one level through each of them, for the case of S^2 phase space. The band topology is described by two integers (first and second Chern classes) which play the role of the topological quantum numbers for

³ To be precise, $Ch(Polyad) = \exp(Nx)$ is the Chern Character of a holomorphic line bundle which enters in the geometric quantization description of the space $\mathcal{H}_{polyad}$. See [25] chap.8.

⁴The value $\Delta N = (N + 1) + 1$ can be interpreted as $(N + 1)$ quantized states of the degeneracy sphere plus an extra level due to the topology of the degeneracy bundle.

energy bands. Topologically coupled energy bands (i.e. non decomposable vector bundles), appear for problems with four-dimensional $\mathbb{C}P^2$ phase space, whereas for S^2 phase space one needs only a single Chern index to characterize the coupling with fast quantum subsystem, and consequently every elementary band has rank 1.

It should be noted that the coupling of a slow classical subsystem with at least three electronic quantum states is a necessary condition to see the formation of topologically non trivial bands from initially isolated trivial bands. In the above model, two isolated trivial bands can not be transformed into non-trivial bands for topological reasons [27]. This is just because the Chern class of a trivial rank 2 bundle over $\mathbb{C}P^2$ is $C(T) = 1$, and the decomposition into two rank one bundles $T = F_1 \oplus F_2$ gives the factorization $C(\mathbb{C}^2) = 1 = (1 + A_1x)(1 + A_2x)$ which implies that $A_1 = A_2 = 0$, and that F_1 and F_2 are trivial bundles. This constraint does not exist in the case of vector bundles over sphere S_2 , where the Chern class is just a degree one polynomials (one just has then $A_1 = -A_2$).

This precedent remark does not mean that two isolated trivial bands cannot become coupled due to vibronic interaction: the formation of degeneracy surfaces between two bands is naturally possible.

4 Conclusions

The model presented in this paper can be adapted to different situations in molecular physics where the slow motion phase space can be $S^2, \mathbb{C}P^2, \mathbb{C}P^3, \dots$ and products of them, depending of the relevant degrees of freedom, which can be rotational, vibrational, or electronic. As a concrete molecular object which fits well the assumptions made by the present model, we can mention the ro-vibronic structure of Jahn-Teller molecule $V(CO)_6$ [22]. This concrete example needs a slight extension to the model with a $\mathbb{C}P^2 \times S^2$ slow phase space coupled with three quantum states, and is especially interesting for further analysis.

Our approach can also be used in other areas of physics where Born-Oppenheimer approximation is relevant and where the slow subsystem can be considered as classical and coupled to several quantum states.

>From the mathematical point of view we have shown a concrete manifestation of Atiyah-Singer's index formula in molecular physics. This gives new field of

applications to the index formula. Up to now it was mainly related to quantum chromodynamics with the topology of instantons and gluon fields, but for different reasons ([24] p.355). The topology of bands are also known to have an importance in solid state physics for the integer quantum Hall effect [28, 29]. Molecular systems supply a lot of new examples where the qualitative and quantitative predictions of Atiyah-Singer's index formula can be unambiguously tested and verified with high accuracy experimental measurements and numerical calculations.

Acknowledgements The work is supported by the EU project Mechanics and Symmetry in Europe contract #HPRN-CT-2000-00113.

References

- [1] L. Michel and B. I. Zhilinskii, Phys. Rep. **341** (2001) 11.
- [2] B. I. Zhilinskii, Phys. Rep. **341** (2001) 85.
- [3] W. Harter, Comp. Phys. Rep. **8** (1988) 319.
- [4] H. Cycon, R. Froese, W. Kirsch, and B. Simon, Schrödinger Operators, Springer-Verlag, 1987.
- [5] M. Kellman, J. Chem. Phys. **93** (1990) 6630.
- [6] D. Sadovskii, B. Zhilinskii, J. Chem. Phys. **103** (1995) 10520.
- [7] V.B. Pavlov-Verevkin, D.A. Sadovskii, and B. I. Zhilinskii, Europhys. Lett. **6** (1988) 573;
- [8] B.I. Zhilinskii and S. Brodersen, J. Mol. Spectrosc. **163** (1994) 326;
- [9] S. Brodersen and B.I. Zhilinskii, J. Mol. Spectrosc. **169** (1995) 1;
- [10] D.A. Sadovskii, and B.I. Zhilinskii, Phys. Lett. A, **256** (1999) 235;
- [11] L.Grondin, D.A. Sadovskii, and B.I. Zhilinskii, Phys. Rev. A, **65** (2002).
- [12] F. Faure and B. Zhilinskii, Phys. Rev.Lett. **85** (2000) 960.
- [13] F. Faure and B. Zhilinskii, Lett. Math. Phys. **55**, (2001) 219.

- [14] E. Hawkins, Commun. Math. Phys. **215** (2000) 409.
- [15] B. Fedosov, Commun. Math. Phys. **209** (2000) 691.
- [16] Emmrich, C. and Weinstein, A., Commun. Math. Phys. **176** (1996) 701.
- [17] B. Zhilinskii, Chem. Phys. **137** (1989) 1.
- [18] R. Englman, *The Jahn-Teller effect in molecules and crystals*. Wiley, New-York, (1972).
- [19] J. Hougen. In Physical Chemistry. Advanced Treatise. Vol 4, N.Y. 1969.
- [20] B. R. Judd. Can. J. Phys. **52** (1974) 999.
- [21] M. Rey, V. Boudon, M. Loëte, F. Michelot. J. Mol. Spectrosc. **204** (2000) 106;
- [22] M. Rey, V. Boudon, M. Loëte, P. Asselin, P. Soulard, L. Manceron. J. Chem. Phys. **114** (2001) 10773.
- [23] L.D. Landau and E.M. Lifshitz, *Quantum Mechanics. Non relativistic theory*. Addison-Wesley (1958).
- [24] T. Eguchi, P. Gilkey, and A. Hanson, Phys. Rep. **66** (1980) 213.
- [25] N.M.J. Woodhouse, *Geometric Quantization*, Clarendon Press Oxford (1992).
- [26] J. J. Duistermaat, G. J. Heckman. Invent. Math. **69** (1982) 259.
- [27] F. Faure, B. I. Zhilinskii. Acta Appl. Math. (2001), in press.
- [28] D. J. Thouless, *Topological quantum numbers in non relativistic physics*. (World Scientific, Singapore), 1998.
- [29] F. Faure, J. Phys. A, Math. Gen. **33** (2000) 531.